

New percolation crossing formulas and second-order modular forms

Nikolaos Diamantis

School of Mathematical Sciences

University of Nottingham

Nottingham, UK

and

Peter Kleban

LASST and Department of Physics & Astronomy

University of Maine, Orono, Maine, USA

May 11, 2009

Abstract

We consider the three new crossing probabilities for percolation recently found via conformal field theory [23]. We prove that all three of them (i) may be simply expressed in terms of Cardy's [4] and Watts' [24] crossing probabilities, (ii) are (weakly holomorphic) second-order modular forms of weight 0 (and a single particular type) on the congruence group $\Gamma(2)$, and (iii) under some technical assumptions (similar to those used in [19]), are completely determined by their transformation laws.

The only physical input in (iii) is Cardy's crossing formula, which suggests an unknown connection between all crossing-type formulas.

1 Introduction

Modular behavior appears in percolation theory on a rectangle, as is demonstrated in [19]. There, formulas for the crossing probabilities at the percolation point obtained via conformal field theory (or more exactly, their derivatives with respect to the aspect ratio r) are shown to have interesting modular properties. Further, a modular characterization of these probabilities is given via several theorems.

In that work, there are two main quantities of interest. The “horizontal” crossing probability $\Pi_h(r)$ is the probability of finding at least one cluster that connects the left and right (vertical) edges of the rectangle, with r being the ratio of horizontal to vertical edge lengths. An explicit formula for this quantity was originally calculated by Cardy [4] via conformal field theory, and more recently proven rigorously, for the triangular lattice, by Smirnov [22]. The “horizontal but not vertical” crossing probability $\Pi_{h\bar{v}}(r)$ is the probability of finding at least one cluster that connects the left and right edges of

the rectangle while the top and bottom (horizontal) edges are not connected. A formula for this was first obtained with conformal field theory by Watts [24] and recently proven rigorously by Dubédat [9] using Schramm-Loewner Evolution methods.

The behavior of the crossing probabilities under $S : z \rightarrow -1/z$ (with $z = ir$) follows from physical symmetries of the problem, while the behavior under $T : z \rightarrow z + 1$ arises from the structure of the formulas themselves (explicitly, the appearance of a single “conformal block”, see [19] for details), but has no obvious physical origin.

Two surprises arise in [19]. First, there is no reason to expect modular behavior on a rectangle at all since it lacks the appropriate symmetry. (By contrast, on a torus modular behavior of crossing probabilities is both expected and observed ([20])). Despite this, $\Pi'_h(r)$ is a modular form (the prime denotes differentiation), and is completely determined by a simple modular argument that assumes its physical symmetry. Secondly, $\Pi'_{h\bar{v}}(r)$ is observed to possess unusual modular behavior, which leads to the definition of a new kind of modular object, the n^{th} – *order modular form*. Such objects were also defined independently, in a different context ([8]) and have since been explored systematically (e.g. [18], [7], [14], [13], [10]).

For percolation, the appearance of a second-order modular form may be traced to the difference in sign of the behavior under S of Π'_h and Π'_{hv} , where $\Pi_{hv}(r) = \Pi_h(r) - \Pi_{h\bar{v}}(r)$ is the probability of finding a cluster that crosses both horizontally and vertically.

Recently, three new percolation crossing probabilities have been calculated from conformal field theory, $\pi_h^b(\alpha, \beta)$, $\pi_h^{\bar{b}}(\alpha, \beta)$, and $\nu_h(\alpha, \beta)$ ([23]). These quantities (see below for the explicit formulas) are more precisely probability densities for clusters that connect to two specified points on opposite vertical edges of a rectangle, under certain specified conditions (see [23] for details, but note that the points on the rectangle are the images of α and β , which lie on the real axis). Their integrals give Π_h , $\Pi_{h\bar{v}}$, and $\mathcal{N}_h$, where the latter is the expected number of horizontally crossing clusters. (This was calculated first via conformal field theory [5, 6] and later rigorously [17].)

In this work we examine the modular properties of π_h^b , $\pi_h^{\bar{b}}$, and ν_h under the congruence group $\Gamma(2)$. Unexpectedly, we find that all three of them are second-order modular forms. This occurs even though they again are on a rectangle, which does not possess the requisite symmetry under S , and they specify crossings between points, rather than intervals. These functions are also very interesting from a purely number-theoretic viewpoint, as they illustrate the occurrence of an extension of higher order forms which involves *two* group actions and whose study, though formally natural, would seem otherwise unmotivated.

Proposition 2.2 shows that all three crossing probabilities may be written as expressions linear in the ratio ϕ , which is proportional to $\Pi'_{h\bar{v}}/\Pi'_h$ (see (11), the prime denotes a derivative), with coefficients that are algebraic functions of the cross-ratio λ of the four points defining the rectangle.

Theorem 3.1 proves that all three crossing probabilities are weakly holomorphic second-order modular forms (definitions are given in Section 3), and determines their leading Fourier terms at each cusp. Weak holomorphicity has been studied and led to striking results including the resolution of classical number-theoretical problems on partitions of integers and other results by D. Zagier, R. Borcherds, K. Ono, K. Bringmann ([1, 25, 2, 3], see also [11] for further references). However we did not expect it in this context.

Theorem 4.5 is a Hamburger-type result for the weakly holomorphic higher-order

forms studied in this work. It proves that, under some technical assumptions analogous to those used in [19], all three crossing probabilities are determined by the transformation laws they satisfy. Since the only physical input to the theorem is Π'_h , one finds that a single basic framework of assumptions leads to all three probability functions. This points to an unknown connection between the functions Π_h , $\Pi_{h\bar{v}}$, and $\mathcal{N}_h$.

This work is focused on the mathematical aspects of the new crossing probabilities. We plan to explore its physical implications more fully elsewhere. However, since the treatment here may also be of interest to physicists, we have included some comments that are intended to make the presentation more accessible to that audience.

2 Crossing probabilities

Set $T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. We let $\Gamma(2)$ denote the group of matrices of $\text{SL}_2(\mathbb{Z})$ congruent to the identity mod(2). We use the set of generators of $\Gamma(2)$ consisting of $g_1 := T^2$ and $g_2 := ST^{-2}S^{-1} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$ ([21], p. 63).

Now set $q := e^{2\pi iz}$. Let η be the Dedekind eta function

$$\eta(z) = q^{1/24} \prod_{n=1}^{\infty} (1 - q^n).$$

Also, let λ denote the classical modular function for $\Gamma(2)$. It is given by

$$\lambda(z) := 16 \frac{\eta(z/2)^8 \eta(2z)^{16}}{\eta(z)^{24}} = 1 - \frac{\eta(z/2)^{16} \eta(2z)^8}{\eta(z)^{24}}. \quad (1)$$

Neither the function λ nor its derivative with respect to z

$$\lambda'(z) = 16\pi i \frac{\eta(z/2)^{16} \eta(2z)^{16}}{\eta(z)^{28}} \quad (2)$$

have any poles or zeros in $\mathfrak{H}$ (cf. [21], Section 7.2, (7.2.17)).

$\lambda(z)$ appears here because for $z = ir$ it is the cross-ratio of the four points to which the corners of a rectangle of aspect ratio r are mapped.

Now let

$$C := \frac{2^{1/3} \pi^2}{3 \Gamma(1/3)^3}, \quad (3)$$

following the notation in [19]. Then the horizontal crossing probability $\Pi_h(\lambda(ir))$ satisfies

$$\frac{d}{dr} (\Pi_h(\lambda(ir))) = -4\sqrt{3} C \eta(ir)^4, \quad (4)$$

as proven in [26].

The horizontal but not vertical crossing probability $\Pi_{h\bar{v}}(\lambda(ir))$ is proven in [19] to satisfy

$$\frac{d}{dr} (\Pi_{h\bar{v}}(\lambda(ir))) = -8\sqrt{3} f_2(ir), \quad (5)$$

where

$$\begin{aligned} f_2(z) &= \frac{2\pi i}{3} \eta(z)^4 \int_{\infty}^z f_3(w) dw, \text{ with} \\ f_3(w) &:= \frac{\eta(w/2)^8 \eta(2w)^8}{\eta(w)^{12}}. \end{aligned} \quad (6)$$

The Dedekind eta function is a weight $1/2$ cusp form on $\mathrm{SL}_2(\mathbb{Z})$ with a character. We denote the character of $\eta(z)^4$ by χ . As shown in [19], $f_2(z)$ is a second-order modular form, of a kind which we specify in the next section.

Now let

$$g(z) := \eta(z/2)^8 \eta(2z)^8 / \eta(z)^{16}. \quad (7)$$

This definition is useful for avoiding ambiguities from fractional powers of $\lambda(1 - \lambda)$. Because of (1) g satisfies

$$g^3 = \frac{1}{16} \lambda(1 - \lambda). \quad (8)$$

Then, combining (1) and (2) gives, for $z \in \mathfrak{H}$,

$$\eta(z)^4 = \frac{1}{16\pi i} \frac{\lambda'(z)}{g(z)^2}, \quad (9)$$

as well as

$$f_3(z) = \frac{1}{16\pi i} \frac{\lambda'(z)}{g(z)}. \quad (10)$$

Now let

$$\phi(z) := \frac{f_2(z)}{\eta(z)^4} = \frac{1}{2} C \frac{\Pi'_{h\bar{v}}(z)}{\Pi'_h(z)}. \quad (11)$$

Since $\Pi_{hv}(i/r) = \Pi_{hv}(ir)$ by symmetry, $\Pi'_{hv}(i) = 0$ (note that $r = 1$ is a square). Further, $\Pi_{hv} = \Pi_h - \Pi_{h\bar{v}}$, so that $\Pi'_h(i) = \Pi'_{h\bar{v}}(i)$. Thus

$$\phi(i) = \frac{C}{2}. \quad (12)$$

Using (6) to calculate ϕ' and (10), one sees that ϕ is a function of λ only. Integrating gives

$$\phi(z) = \frac{1}{2^{8/3}} \lambda(z)^{2/3} {}_2F_1(1/3, 2/3; 5/3; \lambda(z)). \quad (13)$$

In a more recent work ([23]), the three new crossing probabilities mentioned in the Introduction are introduced and computed. For $0 < \alpha < 1$, and $1 < \beta$, they are given by

$$\pi_h^b(\alpha, \beta) = \frac{(\beta + \alpha) {}_2F_1(1, 4/3, 5/3, 1 - \alpha/\beta) - 2\beta}{4\sqrt{3} \pi \beta^2 (\beta - \alpha)}, \quad (14)$$

$$\pi_h^{\bar{b}}(\alpha, \beta) = \frac{(\beta + \alpha) {}_2F_1(1, 4/3, 5/3, \alpha/\beta) + 2\beta}{4\sqrt{3} \pi \beta^2 (\beta - \alpha)}, \text{ and} \quad (15)$$

$$\nu_h(\alpha, \beta) = \frac{\beta^2 + 2\alpha\beta - (\beta^2 - \alpha^2) {}_2F_1(1, 4/3, 5/3, \alpha/\beta)}{4\sqrt{3} \pi \beta^2 (\beta - \alpha)^2}. \quad (16)$$

In [23] it is shown that a double integration of each of these quantities gives the crossing probabilities studied in [19] as well as $\mathcal{N}_h$. Specifically

$$\frac{1}{2} \Pi_{h\bar{v}}(\lambda) = \int_0^\lambda \int_1^\infty \pi_h^{\bar{b}}(\alpha, \beta) d\beta d\alpha, \quad (17)$$

$$\Pi_h(\lambda) - \frac{1}{2} \Pi_{h\bar{v}}(\lambda), = \int_0^\lambda \int_1^\infty \pi_h^b(\alpha, \beta) d\beta d\alpha, \text{ and} \quad (18)$$

$$-\frac{1}{2} \Pi_{h\bar{v}}(\lambda) + \frac{\sqrt{3}}{4\pi} \log\left(\frac{1}{1 - \lambda}\right) = \int_0^\lambda \int_1^\infty \nu_h(\alpha, \beta) d\beta d\alpha. \quad (19)$$

(The lhs of (19) is equal to $\mathcal{N}_h(\lambda) - \Pi_h(\lambda)$).

For clarity, we now introduce the notation

$$\begin{aligned} p_{\bar{b}}(z) &:= \pi_{\bar{h}}^{\bar{b}}(\lambda(z), 1), \\ p_b(z) &:= \pi_h^b(\lambda(z), 1), \text{ and} \\ n(z) &:= \nu_h(\lambda(z), 1), \end{aligned} \tag{20}$$

for the holomorphic functions of $z \in \mathfrak{H}$ obtained from (14), (15), and (16) after we replace α and β with $\lambda(z)$ and 1 respectively.

We next prove (with prime standing for differentiation in z)

Theorem 2.1. *For all $z \in \mathfrak{H}$,*

$$\lambda'(z) p_{\bar{b}}(z) = 4\sqrt{3}i \left(\frac{\lambda(z)f_2(z)}{\lambda'(z)} \right)', \tag{21}$$

$$\lambda'(z) p_b(z) = 4\sqrt{3}Ci \left(\frac{\lambda(z)}{\lambda'(z)} \eta^4 \right)' - 4\sqrt{3}i \left(\frac{\lambda(z)f_2(z)}{\lambda'(z)} \right)', \text{ and} \tag{22}$$

$$\lambda'(z) n(z) = \frac{\sqrt{3}}{4\pi} \left(\frac{\lambda(z)}{1 - \lambda(z)} \right)' - 4\sqrt{3}i \left(\frac{\lambda(z)f_2(z)}{\lambda'(z)} \right)'. \tag{23}$$

Proof. We first deduce from (15) that $\pi_{\bar{h}}^{\bar{b}}(\alpha, \beta) = \beta^{-2} \pi_{\bar{h}}^{\bar{b}}(\alpha/\beta, 1)$. This and the change of variables $\alpha/\beta \rightarrow \beta$ imply that the rhs of (17) equals

$$\int_0^\lambda \int_0^\alpha \pi_{\bar{h}}^{\bar{b}}(x, 1) dx \frac{d\alpha}{\alpha}$$

and hence a differentiation in terms of ir together with (5) (recalling that the differentiation in (5) is in terms of r and not ir) and re-arranging gives

$$4i\sqrt{3} \frac{\lambda(ir)f_2(ir)}{\lambda'(ir)} = \int_0^{\lambda(ir)} \pi_{\bar{h}}^{\bar{b}}(x, 1) dx.$$

One more differentiation in terms of ir and the definitions (20) implies the result for $z = ir$. The analytic continuation of $\pi_{\bar{h}}^{\bar{b}}(\cdot, 1)$ and the fact that λ, λ' do not have any poles or zeros in $\mathfrak{H}$ imply that it holds for all $z \in \mathfrak{H}$.

The assertions about the other two crossing probabilities are proved in exactly the same way using (18), (19). $\square$

We next re-write $p_{\bar{b}}$, p_b , and n in a way that avoids derivatives of λ

Proposition 2.2. *For every $z \in \mathfrak{H}$, we have*

$$\begin{aligned} p_{\bar{b}}(z) &= \frac{1}{4\sqrt{3}\pi g(z)^2} \frac{1 + \lambda(z)}{1 - \lambda(z)} \phi(z) + \frac{1}{2\sqrt{3}\pi} \frac{1}{1 - \lambda(z)} \\ p_b(z) &= \frac{1}{4\sqrt{3}\pi g(z)^2} \frac{1 + \lambda(z)}{1 - \lambda(z)} (2\phi(i) - \phi(z)) - \frac{1}{2\sqrt{3}\pi} \frac{1}{1 - \lambda(z)} \\ n(z) &= -\frac{1}{4\sqrt{3}\pi g(z)^2} \frac{1 + \lambda(z)}{1 - \lambda(z)} \phi(z) + \frac{1}{4\sqrt{3}\pi} \frac{1 + 2\lambda(z)}{(1 - \lambda(z))^2}. \end{aligned}$$

Proof. By (10),

$$\lambda'(z) = 16\pi i g(z)^2 \eta(z)^4. \quad (24)$$

Using (8) and (24),

$$3g'g^2 = \frac{1}{16}(1-2\lambda)\lambda' = \pi i(1-2\lambda)g^2\eta^4.$$

This and (24) imply that

$$\left(\frac{\lambda}{\lambda'}\eta^4\right)' = \left(\frac{\lambda}{16\pi i g^2}\right)' = \frac{1}{16\pi i} \frac{\lambda'g - 2g'\lambda}{g^3} = \frac{\eta^4(1+\lambda)}{3(1-\lambda)}$$

Expanding (21) according to this formula, from (10) and the definition of f_2 we deduce

$$p_{\bar{b}}(z) = \frac{1}{4\sqrt{3}\pi g(z)^2} \frac{1+\lambda(z)}{1-\lambda(z)} \phi(z) + \frac{1}{2\sqrt{3}\pi} \frac{1}{1-\lambda(z)}.$$

The proof of the formula for p_b is similar. The formula for n may be obtained by comparing (15) with (16) and making use of the formula for $p_{\bar{b}}$. $\square$

Remark 2.3. Using (8) to write g in terms of λ , Proposition 2.2 gives each of $p_{\bar{b}}$, p_b , and n as an expression linear in the ratio ϕ (see (11)), with the coefficients being rational functions of λ . Thus all three probabilities are determined by the same physical quantity ϕ , which is itself proportional to $\Pi'_{h\bar{v}}/\Pi'_h$. Note also that the constant $C = 2\phi(i)$ in p_b is exactly the constant C in (39) of [19]. It follows that $2\phi(i) - \phi(z)$ and $\phi(z)$ transform into each other under S .

Remark 2.4. Proposition 2.2 illustrates why it is natural to work with the group $\Gamma(2)$ rather than the *theta group* Γ_θ , as in [19]. The ratio ϕ is easily seen—use, e.g. (39) of [19]—to transform as a second-order modular form (for the definition see [19] or the next section) under the generators of either $\Gamma(2)$ (e.g. $\{g_1, g_2\}$) or Γ_θ (e.g. $\{g_1, S\}$). However, $\lambda|_0 S = 1 - \lambda$; as a result the rational functions of λ of Remark 2.3 do not transform simply under Γ_θ .

3 Higher-order forms

Let Γ be a congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$ and $k \in 2\mathbb{Z}$. As usual, for every character χ on Γ we define an action of Γ on the space of functions $f : \mathfrak{H} \rightarrow \mathbb{C}$ given, for $\gamma \in \Gamma$, by

$$(f|_{k,\chi}\gamma)(z) = f(\gamma z) j(\gamma, z)^{-k} \overline{\chi(\gamma)}.$$

Here, $j\left(\begin{pmatrix} * & * \\ c & d \end{pmatrix}\right) := cz + d$. When χ is the trivial character $\mathbf{1}$ we write $f|_k \gamma$.

To describe the condition on the “growth at the cusps” which will be included in the definition of higher-order forms we first note that a *cusp* is a point $x \in \mathbb{R} \cup \{\infty\}$ such that $\gamma x = x$ for some $\pm 1 \neq \gamma \in \Gamma$ with $|\mathrm{tr}(\gamma)| = 2$. Cusps are important because they are the only points on the real line at which modular forms have a regular behavior. Indeed, one of their special features is that if a function is Γ -invariant under the action $|_k$, then its composition by an appropriate map sending infinity to a cusp will be periodic and hence it will possess a Fourier expansion. This fact is crucial in establishing the finite dimensionality of the space of modular forms. As a result (as will be seen in the proof of

Theorem 4.2) in order to uniquely determine functions, in addition to the transformation laws satisfied, detailed knowledge of the growth at the cusps is essential.

Two cusps $\mathfrak{a}, \mathfrak{b}$ are called *equivalent* if there is a $\gamma \in \Gamma$ such that $\mathfrak{a} = \gamma\mathfrak{b}$ and *inequivalent* otherwise.

For each cusp $\mathfrak{a}$ there is a *scaling matrix*, i.e. a matrix $\sigma_{\mathfrak{a}}$ such that

$$\sigma_{\mathfrak{a}}(\infty) = \mathfrak{a} \quad \text{and} \quad \sigma_{\mathfrak{a}}^{-1} \Gamma_{\mathfrak{a}} \sigma_{\mathfrak{a}} = \Gamma_{\infty}$$

where $\Gamma_{\mathfrak{a}}$ (resp. Γ_{∞}) is the set of elements of Γ fixing $\mathfrak{a}$ (resp. ∞) ([12], Section 2). If f is a function on $\mathfrak{H}$ such that $f|_k \sigma_{\mathfrak{a}}$ has a Fourier expansion of the form

$$(f|_k \sigma_{\mathfrak{a}})(z) = \sum_{n=-m}^{\infty} a_n e^{\pi i n z / a}$$

for some $m \in \mathbb{Z}_{\geq 0}$ with $a \in \mathbb{Q}_+$ and $a_{-m} \neq 0$, then we say that f is *meromorphic* at $\mathfrak{a}$. If $m = 0$ (resp. $m < 0$), then f is *holomorphic* (resp. *cuspidal*) at $\mathfrak{a}$.

For a set $\{\chi_1, \dots, \chi_n\}$ of characters on Γ , we define a *weakly holomorphic n th-order modular form on Γ of weight k and type $(\chi_1, \dots, \chi_n)$* to be a holomorphic function on $\mathfrak{H}$ which is meromorphic at the cusps and such that, for every $\gamma_i \in \Gamma$,

$$f|_{k, \chi_1}(\gamma_1 - 1)|_{k, \chi_2}(\gamma_2 - 1) \dots |_{k, \chi_n}(\gamma_n - 1) = 0.$$

If f is holomorphic (resp. cuspidal) at all cusps, then we call the form holomorphic (resp. cuspidal). We denote the space of order n weakly holomorphic (resp. holomorphic, cuspidal) forms of weight k and type $(\chi_1, \dots, \chi_n)$ on Γ by $\tilde{M}_k^n(\Gamma; \chi_1, \dots, \chi_n)$ (resp. $M_k^n(\Gamma; \chi_1, \dots, \chi_n)$, $S_k^n(\Gamma; \chi_1, \dots, \chi_n)$).

Note that, in contrast to the case of ordinary modular forms, a Fourier expansion at the cusps is not guaranteed. The reason is that this would necessitate that f be invariant under a subgroup of $\sigma_{\mathfrak{a}} \Gamma_{\infty} \sigma_{\mathfrak{a}}^{-1}$ which, in general, does not happen when the order is higher than 1.

We next prove that the three new crossing probabilities $p_{\bar{b}}$, p_b , and n are second-order forms and determine their growth at the cusps.

The group $\Gamma(2)$ has three inequivalent cusps at ∞ , 0 and -1 . Three corresponding scaling matrices are I , $U := \begin{pmatrix} 0 & -1 \\ 1 & 1 \end{pmatrix} = ST$ and $U^2 = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$.

We can then prove

Theorem 3.1. *As functions of z , $p_{\bar{b}}(z)$, $p_b(z)$ and $n(z)$ are weakly holomorphic second-order modular forms on $\Gamma(2)$ of weight 0 and type $(\mathbf{1}, \chi)$. The first power of q appearing in the expansion of $p_{\bar{b}}$ at ∞ (resp. 0 , -1) is 1 (resp. $q^{-5/6}$, $q^{2/3}$). The corresponding first powers for p_b and n are $q^{-1/3}$, 1 , $q^{2/3}$ and $q^{1/2}$, q^{-1} , $q^{2/3}$.*

Proof. The function f_2 is a holomorphic weight 2 second-order form of type $(\mathbf{1}, \chi)$ satisfying

$$f_2|_2(\gamma - 1) = d_{\gamma} \eta^4 \quad \text{for all } \gamma \in \Gamma_{\theta} \quad (25)$$

for a constant d_{γ} depending solely on γ ([19], (38)). Since $\Gamma(2)$ is a subgroup of Γ_{θ} , (25) holds for $\Gamma(2)$ too. It is easy to verify that $d_{g_1} = 0$ and $d_{g_2} = C(e^{-2\pi i/3} - 1)$, so the rhs of (25) is not zero for all $\gamma \in \Gamma(2)$.

An easy computation (see also Theorem 2.2 of [7]) implies that, for $\gamma \in \Gamma(2)$,

$$\left(\frac{\lambda}{\lambda'} f_2\right)|_0 \gamma = \left(\frac{\lambda}{\lambda'}\right)|_{-2} \gamma f_2|_2 \gamma = \frac{\lambda}{\lambda'} f_2 + \frac{d_{\gamma} \lambda \eta^4}{\lambda'}.$$

Therefore

$$\left(\frac{\lambda}{\lambda'} f_2\right)'|_2(\gamma-1) = \left(\frac{d_\gamma \lambda \eta^4}{\lambda'}\right)' . \quad (26)$$

which, by Theorem 2.1, implies

$$p_{\bar{b}}|_0(\gamma-1) = d_\gamma G \quad (27)$$

where

$$G := \frac{1}{\lambda'} \left(\frac{\lambda \eta^4}{\lambda'}\right)' .$$

Since $\lambda \eta^4 / \lambda'$ transforms as a modular form of weight 0 with character χ on $\Gamma(2)$, its derivative transforms as a weight 2 form and, therefore, G transforms as a weight 0 form with character χ .

The behaviour of p_b under the action of $\Gamma(2)$ follows from (22), which implies that $p_b = -p_{\bar{b}} + 4\sqrt{3} C i G$. Since $G|_0(\gamma-1) = (\chi(\gamma)-1)G$, we conclude that $p_b|_0(\gamma-1)$ transforms as a weight 0 modular form with character χ .

In the same way, since $(\lambda/(1-\lambda))'/\lambda'$ transforms as a weight 0 modular form with trivial character, we deduce that $n|_0(\gamma-1)$ transforms as a weight 0 modular form with character χ .

To complete the proof that $p_{\bar{b}}(z)$, $p_b(z)$ and $n(z)$ are second-order weakly holomorphic forms of weight 0 and type $(\mathbf{1}, \chi)$ for $\Gamma(2)$, it remains to show that $p_{\bar{b}}(z)$ and G are meromorphic at the cusps. The argument also proves the last part of the theorem, specifying the first terms of the Fourier expansions of the three probability functions.

We start by determining the first powers of q appearing in the Fourier expansions of λ at the three inequivalent cusps ∞ , 0 and -1 . We shall be making use of (9), (13), and the transformations of λ : $\lambda|_0 S = 1 - \lambda$ and $\lambda|_0 T = \lambda/(\lambda-1)$ ([21], (7.2.2)). Thus

$$\lambda|_0 U = 1/(1-\lambda) \quad \text{and} \quad \lambda|_0 U^2 = 1 - 1/\lambda. \quad (28)$$

Since the leading term in λ at ∞ is $16q^{1/2}$, it follows that the leading terms at 0 and -1 are $1 + 16q^{1/2}$ and $q^{-1/2}$ respectively. Therefore λ' is a weight 2 first-order weakly holomorphic form on $\Gamma(2)$ and the first powers in its Fourier expansions at the cusps ∞ , 0 and -1 are $q^{1/2}$, $q^{1/2}$ and $q^{-1/2}$ respectively. The corresponding powers for λ/λ' are then 1, $q^{-1/2}$ and 1.

Next we verify that $p_{\bar{b}}(z)$, $p_b(z)$ and $n(z)$ have Fourier expansions at $\infty, 0, -1$. We first note that the Fourier expansion of λ and (28) imply that λ maps the cusps ∞ , 0 and -1 to 0, 1, and ∞ respectively. On the other hand, each of the probabilities $\pi_{\bar{h}}^b(\lambda(z), 1)$, $\pi_h^b(\lambda(z), 1)$ and $\nu_h(\lambda(z), 1)$ is given (see (14), (15), and (16)) in terms of a hypergeometric function of λ or $1-\lambda$ and integral powers of λ only. It follows from the classical linear transformation formulas for hypergeometric functions that an expansion around any of the cusps will involve integral powers of λ or $\lambda^{1/3}$ only. Thus any of these expansions is invariant under T^6 , which establishes the periodicity of g , $g|_0 U$, $g|_0 U^2$, for $g = \pi_{\bar{h}}^b(\lambda(z), 1)$, $\pi_h^b(\lambda(z), 1)$ and $\nu_h(\lambda(z), 1)$.

The leading term in each of $\pi_{\bar{h}}^b(\lambda, 1)$, $\pi_h^b(\lambda, 1)$ and $\nu_h(\lambda, 1)$ and therefore $p_{\bar{b}}(z)$, $p_b(z)$ and $n(z)$ at each cusp then follows immediately from (14), (15) and (16) using the Taylor expansion of the hypergeometric function and $\lambda \sim q^{1/2}$.

The first terms of the Fourier expansions of G follow in similar (but simpler) way from the Fourier expansions of λ , λ' and the observation that the first power in the expansion of η^4 is $q^{1/6}$.

Table 1: First terms of Fourier expansions

	∞	0	-1
λ	$q^{1/2}$	$1 + 16q^{1/2}$	$q^{-1/2}$
$\frac{\lambda}{\lambda'}$	1	$q^{-1/2}$	1
λ'	$q^{1/2}$	$q^{1/2}$	$q^{-1/2}$
$\frac{1}{\lambda'}$	$q^{-1/2}$	$q^{-1/2}$	$q^{1/2}$
$p_{\bar{b}}(z)$	1	$q^{-5/6}$	$q^{2/3}$
$p_b(z)$	$q^{-1/3}$	1	$q^{2/3}$
$n(z)$	$q^{1/2}$	q^{-1}	$q^{2/3}$
$\left(\frac{\lambda}{\lambda'}\eta^4\right)'$	$q^{1/6}$	$q^{-1/3}$	$q^{1/6}$
$\frac{1}{\lambda'}\left(\frac{\lambda}{\lambda'}\eta^4\right)'$	$q^{-1/3}$	$q^{-5/6}$	$q^{2/3}$

Finally, one must also verify the meromorphicity at all equivalent cusps. For G , this is automatically implied by its invariance under the action of $\Gamma(2)$. For the three probability functions, it is no longer automatic because they are not $\Gamma(2)$ -invariant. Let $\mathfrak{b} = \gamma \mathfrak{a}$ ($\gamma \in \Gamma(2)$) for $\mathfrak{a} = \infty, 0$ or -1 . We have $\sigma_{\mathfrak{b}} = \gamma \sigma_{\mathfrak{a}}$, $\Gamma_{\sigma_{\mathfrak{b}}} = \gamma \Gamma_{\sigma_{\mathfrak{a}}} \gamma^{-1}$ and that if $\sigma_{\mathfrak{b}}'$ is another scaling matrix for $\mathfrak{b}$ then $\sigma_{\mathfrak{b}} = \sigma_{\mathfrak{b}}' T^m$, with $m \in 2\mathbb{Z}_{\geq 0}$ ([12], Section 2). Therefore, if $\sigma_{\mathfrak{b}}'$ is any scaling matrix of $\mathfrak{b}$, we have (with (27))

$$p_{\bar{b}}|_0 \sigma_{\mathfrak{b}}' = p_{\bar{b}}|_0 \sigma_{\mathfrak{b}} T^m = p_{\bar{b}}|_0 \gamma \sigma_{\mathfrak{a}} T^m = p_{\bar{b}}|_0 \sigma_{\mathfrak{a}} T^m + c_{\gamma} G|_0 \sigma_{\mathfrak{a}} T^m$$

From the behaviour of $p_{\bar{b}}$ at $\infty, 0, -1$ we proved above as well as the behaviour of G at these cusps, we deduce the meromorphicity of $p_{\bar{b}}$ at all cusps $\mathfrak{b}$. The same argument applies to p_b and n .

For convenience, Table 1 lists the first terms of the Fourier expansions for the functions discussed.

□

Remark 3.2. Theorem 3.1, in combination with Prop. 2.2, shows that ϕ is a weight 0 weakly holomorphic second-order modular form on $\Gamma(2)$ of type $(\mathbf{1}, \bar{\chi})$. This result is already implied in [19], but here it is proved explicitly.

4 Uniqueness theorems

We call *conformal block* of dimension $\alpha \in \mathbb{R}$ a function of $r > 0$ expressible in the form

$$\sum_{n=0}^{\infty} a_n e^{-\pi(n+\alpha)r}, \quad (29)$$

with $a_0 \neq 0$. As noted in [19], if $\Pi(r)$ is a conformal block, then the convergence of (29) implies that the function

$$P(ir) := \Pi(r), \quad (30)$$

extends to a holomorphic function $P(z)$ with $z \in \mathfrak{H}$ (its power series expansion is exactly as in (29), except that $-r$ is replaced by iz). In [19], transformation properties under S , which maps the imaginary axis into itself, were important. Here, the corresponding group element is $g_2 : z \rightarrow z/(1+2z)$, which does not have this property. Therefore we use a slightly different approach, working exclusively with the analytic continuations of the conformal blocks.

We start with a Lemma which will be needed in the proof of the next theorem.

Lemma 4.1. *A function $f \in S_4(\Gamma(2), \chi)$ is uniquely determined by the coefficients of $e^{\pi iz/3}$ in its Fourier expansions at ∞ and 0.*

Proof of Lemma: For each cusp $\mathfrak{a}$, let

$$E_{\mathfrak{a}}(z) := 2i \lim_{s \rightarrow 1} \frac{d}{dz} \left(\sum_{\gamma \in \Gamma_{\mathfrak{a}} \backslash \Gamma} \text{Im}(\sigma_{\mathfrak{a}} \gamma z)^s \right).$$

Then

$$E_{\mathfrak{a}}(z) - E_{\infty}(z) = -1 - 4\pi^2 \sum_{n=1}^{\infty} n (Z_{\mathfrak{a},\infty}(n, 0, 1) - Z_{\infty,\infty}(n, 0, 1)) e^{2\pi i n z},$$

where $Z_{\mathfrak{a},\mathfrak{b}}(m, n, s)$ is the Selberg-Kloosterman zeta function. (Its definition is given in [15], but is rather lengthy. Since we do not use it further, we refrain from quoting it here.) The set $\{E_{-1} - E_{\infty}, E_0 - E_{\infty}\}$ spans the direct complement of the space of cusp forms within the space of all modular forms of weight 2 for $\Gamma(2)$ ([15]). Further, the dimension of $S_4(\Gamma(2), \chi)$ is 2 ([16], Section 4.). Therefore a basis is given by $(E_0 - E_{\infty})\eta^4$ and $(E_{-1} - E_{\infty})\eta^4$. As a result, any function $f \in S_4(\Gamma(2), \chi)$ has coefficients of $e^{\pi iz/3}$ in its Fourier expansion at ∞ and 0, which we denote by a and b , respectively.

Thus

$$f = x_1(E_0 - E_{\infty})\eta^4 + x_2(E_{-1} - E_{\infty})\eta^4,$$

for some $x_1, x_2 \in \mathbb{C}$. Further (see [15]), for any cusps $\mathfrak{a}, \mathfrak{b}, \mathfrak{c}$,

$$j(\sigma_{\mathfrak{c}}, z)^{-2} (E_{\mathfrak{a}}(\sigma_{\mathfrak{c}} z) - E_{\mathfrak{b}}(\sigma_{\mathfrak{c}} z)) = \delta_{\mathfrak{ac}} - \delta_{\mathfrak{bc}} + O(e^{-2\pi y}),$$

where $z = x + iy$, which gives

$$f = -(x_1 + x_2)e^{\pi iz/3} + O(e^{-7\pi y/3}) \quad f|_4 U = x_1 e^{\pi iz/3} + O(e^{-7\pi y/3}).$$

Thus $x_1 = b$ and $x_2 = -a - b$. □

We are now in a position to prove

Theorem 4.2. *Let $F(z) = \sum_{n=0}^{\infty} b_n e^{\pi i(n+\frac{1}{3})z}$ be the analytic continuation of a conformal block of dimension $1/3$, with $b_0 = \pi i/3$. Suppose that*

$$(a) \quad F|_4 g_2 = \chi(g_2)F, \tag{31}$$

along some curve in $\mathfrak{H}$,

(b) *$F(-1 + i/r)$ is bounded as $r \rightarrow \infty$, and*

(c)

$$\lim_{r \rightarrow \infty} e^{\frac{\pi r}{3}} r^{-4} F\left(\frac{i}{r}\right) = -\frac{4}{3} 2^{1/3} \pi^2. \quad (32)$$

Then $F(z) \in S_4(\Gamma(2), \chi)$ and

$$F(z) = \frac{\lambda'(z)}{\lambda(z)} \left(\frac{\lambda(z) \eta(z)^4}{\lambda'(z)} \right)'.$$

Proof. We first prove that F , if it exists, is a weight 4 cusp form for $\Gamma(2)$ with character χ .

Since F is analytic (31) holds on all of $\mathfrak{H}$. The Fourier expansion of F and $\chi(T^2) = e^{2\pi i/3}$ then imply

$$F|_{4,\chi} T^2 = F.$$

Therefore, $F|_{4,\chi} \gamma = F$, for all $\gamma \in \Gamma(2)$.

We next establish the vanishing at the cusps. First, by its Fourier expansion, F vanishes at ∞ . To verify that the Fourier expansion of $F|_4 U$ has only positive powers (and thus that F vanishes at 0), it suffices to show $(F|_4 U)(-1 + ir) \rightarrow 0$ as $r \rightarrow \infty$. But $(F|_4 U)(-1 + ir) = F(i/r) r^{-4}$, which must vanish as $r \rightarrow \infty$ by (32). For the cusp at -1 consider $(F|_4 U^2)(ir) = F(-1 + i/r) r^{-4}$. By the assumed boundedness of $F(-1 + i/r)$, this vanishes as $r \rightarrow \infty$.

By assumption, the coefficient of $e^{\pi iz/3}$ in the expansion of F at ∞ is

$$b_0 = \lim_{z \rightarrow \infty} e^{-\pi iz/3} F(z) = \frac{\pi i}{3}.$$

The corresponding coefficient at 0 is $\lim_{z \rightarrow \infty} e^{-\pi iz/3} (F|_4 U)(z)$. Taking the limit over the line $z = -1 + ir$, as $r \rightarrow \infty$, and using (32), one has

$$\lim_{r \rightarrow \infty} F(i/r) r^{-4} e^{\pi r/3} e^{\pi i/3} = -\frac{4}{3} 2^{1/3} e^{\pi i/3} \pi^2.$$

A computation confirms that these coincide with the corresponding coefficients for $\frac{\lambda'}{\lambda}(\frac{\lambda}{\lambda'} \eta^4)'$. Lemma 4.1 then implies the result. $\square$

Remark 4.3. Theorem 4.2 resembles Theorem 1 of [19]. In that case, the assumption of modular transformation properties under S led to a proof that an even conformal block (i.e. one with all $a_{2n+1} = 0$) has dimension $1/3$ and is in fact equal to Cardy's function Π_h . Here we assume the block dimension and modular transformation property under the generator g_2 of $\Gamma(2)$ and find that the function is a simple expression involving η^4 , which is proportional to the z -derivative of Π_h . The assumption of evenness is not necessary.

Remark 4.4. Note that if we let $z = ir/(1 + 2ir)$ with $r > 0$ be the curve in (31), the assumption of transformation under g_2 , then the lhs of the equation is in the physical regime, i.e. $z = ir$.

Theorem 4.5. Let $P(z) = \sum_{n=0}^{\infty} a_n e^{\pi i(n+1)z}$ be the analytic continuation of a conformal block of dimension 1, and let $F(z)$ be as in Theorem 4.2. Set

$$P_1 := 4\sqrt{3}C i F - P.$$

Further, for a fixed $A \in \mathbb{C}$ set

$$\tilde{P}(z) := P(z) + A F(z).$$

Suppose that

$$(a) \quad \tilde{P}|_4 g_2 = \tilde{P}, \quad (33)$$

along a curve in $\mathfrak{H}$ and

(b) $P(-1 + i/r)$ is bounded as $r \rightarrow \infty$.

Then

(i) If $A = -C$ and $P(i/r)$ is bounded as $r \rightarrow \infty$,

$$P(z) = \frac{(\lambda'(z))^2}{\lambda(z)} p_{\bar{b}}(z),$$

(ii) Under the same assumptions,

$$P_1(z) = \frac{(\lambda'(z))^2}{\lambda(z)} p_b(z).$$

(iii) If $A = C$ and $P(i/r)r^{-4} \rightarrow -\sqrt{3}\pi/4$ as $r \rightarrow \infty$,

$$P(z) = \frac{(\lambda'(z))^2}{\lambda(z)} n(z).$$

Proof. We consider the claims in order.

Proof of (i) We proceed exactly as in the proof of Theorem 4.2, except replacing χ by 1. Since $\tilde{P}$ (and P) are analytic, if they exist, (33) holds in all of $\mathfrak{H}$. Since $g_2 = ST^{-2}S^{-1}$, (31) and $\chi(T^2) = e^{2\pi i/3}$ give

$$P|_4 (g_2 - 1) = A(1 - e^{-2\pi i/3})F \in S_4(\Gamma(2), \chi). \quad (34)$$

Further, the Fourier expansion of P implies that $P|_4 (T^2 - 1) = 0$. Hence

$$P|_4 (\gamma - 1) \in S_4(\Gamma(2), \chi). \quad (35)$$

As in the proof of Theorem 4.2, we have by assumption that $P(ir)$, $(P|_4 U)(-1 + ir)$, and $(P|_4 U^2)(ir) \rightarrow 0$ as $r \rightarrow \infty$, establishing that $P(z)$ vanishes at all three cusps. By (35), $P|_4 \gamma$ ($\gamma \in \Gamma(2)$) also vanishes at all three cusps.

Now let

$$L := \frac{(\lambda')^2}{\lambda} p_{\bar{b}}.$$

Then

$$\begin{aligned}(P - L)|_{g_1} &= P|_{g_1} - L|_{g_1} \\ &= P - L,\end{aligned}\tag{36}$$

by the periodicity of both P and L . Further

$$(P - L)|_4(g_2 - 1) = A(1 - e^{-2\pi i/3})F - \frac{(\lambda')^2}{\lambda}p_{\bar{b}}|_4(g_2 - 1),$$

by (34). By (27) and Theorem 4.2,

$$p_{\bar{b}}|_4(g_2 - 1) = d_{g_2}G = d_{g_2}\frac{\lambda}{(\lambda')^2}F.$$

Therefore, since $(\lambda')^2/\lambda$ is invariant under g_2 ,

$$\begin{aligned}(P - L)|_4(g_2 - 1) &= A(1 - e^{-2\pi i/3})F - d_{g_2}F \\ &= 0,\end{aligned}\tag{37}$$

using the formula for d_{g_2} after (25). Thus $P - L$ is invariant under both generators of $\Gamma(2)$. As noted when recalling the definition of cusps, this $\Gamma(2)$ -invariance in terms of the action $|_4$ also implies that L has Fourier expansions at each cusp of $\Gamma(2)$. Theorem 3.1 (see Table 1) implies that it vanishes at each cusp. Since P vanishes as well, so does $P - L$, and the latter is a standard weight 4 cusp form with trivial character for $\Gamma(2)$. Therefore it vanishes ([16]), establishing the uniqueness of P (and $\tilde{P}$). This completes the proof of (i).

Proof of (ii): This is immediate from (i), Theorem 4.2, and (22).

Proof of (iii): Analogous to the proof of (i), except that L is redefined with n replacing $p_{\bar{b}}$. Here, neither P nor L vanishes at the cusp at 0, but $P - L$ does, which is sufficient. □

Remark 4.6. Theorem 4.5 resembles Theorem 3 of [19]. In that case, the assumption of modular transformation properties under S for two conformal blocks of arbitrary dimension led to a characterization of crossing probabilities that generalize Π_h and $\Pi_{h\bar{v}}$. Here, by assuming the block dimensions, modular transformation properties under the generator g_2 of $\Gamma(2)$, and cusp properties, we reproduce all three new crossing probabilities.

Remark 4.7. It is interesting that the only input to Theorem 4.5 from the physics of the problem is via F . The physical input to F comes from η^4 , itself proportional to the z -derivative of Π_h . Thus, under the assumptions of the theorem, all three crossing probabilities are determined by Π_h .

5 Acknowledgements

ND thanks Marvin Knopp for useful conversations on certain aspects of the paper. PK is grateful to Don Zagier for leading us to Proposition 2.2, for other useful conversations, and for his kind hospitality at the Collège de France, where part of this research was

done. He also thanks C. Hongler for an advance copy of [17] and J. J. H. Simmons and R. M. Ziff for useful conversations.

This work was supported in part by the National Science Foundation Grant No. DMR-0536927 (PK).

References

- [1] R. Borcherds, *Automorphic forms on $O_{s+2,2}(\mathbb{R})$ and infinite products*, Invent. Math. 120 (1995), no. 1, 161–213.
- [2] K. Bringmann and K. Ono, *The $f(q)$ mock theta function conjecture and partition ranks*, Inv. Math. 165, (2006), 243–266.
- [3] K. Bringmann and K. Ono, *Arithmetic properties of coefficients of half-integral weight Maass-Poincaré series*, Math. Ann. 337 (2007), no. 3, 591–612.
- [4] J. L. Cardy, *Critical percolation in finite geometries*, J. Phys. A **25** 201-206 (1992).
- [5] J. Cardy, *Conformal invariance and percolation*, preprint (2001).
- [6] J. Cardy, *Linking numbers for self-avoiding loops and percolation: Application to the spin quantum Hall transition*, Phys. Rev. Lett. **84** 3507-3510 (2000).
- [7] Y. Choie and N. Diamantis, *Rankin-Cohen brackets on higher-order modular forms*, Proceedings of the Bretton Woods workshop on Multiple Dirichlet series (AMS Proceedings of Symposia in Pure Mathematics).
- [8] G. Chinta, N. Diamantis and C. O’Sullivan, *Second-order modular forms*, Acta Arith. 103 (2002), no. 3, 209–223.
- [9] J. Dubédat, *Excursion decompositions for SLE and Watts’ crossing formula*, Probab. Theory Relat. Fields **134**, 453-488 (2006) (DOI: 10.1007/s00440-005-0446-3).
- [10] A. Deitmar, *Higher order group cohomology and the Eichler-Shimura map*, J. Reine Angew. Math. 629 (2009), 221–235
- [11] W. Duke and P. Jenkins, *Integral traces of singular values of weak Maass forms*, Algebra Number Theory 2 (2008), no. 5, 573–593.
- [12] N. Diamantis, M. Knopp, G. Mason and C. O’Sullivan, *L-functions of second-order cusp forms*, Ramanujan J (2006) 12:327-347.
- [13] N. Diamantis and D. Sim, *The classification of higher-order cusp forms*, J. Reine Angew. Math. 622 (2008), 121–153.
- [14] N. Diamantis and R. Sreekantan, *Iterated integrals and higher order automorphic forms*, Comment. Math. Helv. 81 (2006), no. 2, 481–494.
- [15] D. Goldfeld and C. O’Sullivan, *Estimating additive character sums for Fuchsian groups*, The Ramanujan Journal, 7 (2003), 241–267.
- [16] S. Hussein and M. Knopp, *Eichler cohomology and automorphic forms* Illinois J. Math. 15 1971 565–577.

- [17] C. Hongler and S. Smirnov, *in preparation*.
- [18] Ö. Imamoglu and Y. Martin, *A converse theorem for second-order modular forms of level N* , Acta Arith. 123 (2006), no. 4, 361–376.
- [19] P. Kleban and D. Zagier, *Crossing Probabilities and Modular Forms*, Journal of Statistical Physics, 113 (2003), 431–454.
- [20] H. Pinson, *Critical percolation on the torus*, J. Stat. Phys. **75** 1167–1177 (1994).
- [21] R. A. Rankin, *Modular forms and functions*, Cambridge University Press, Cambridge-New York-Melbourne, 1977.
- [22] S. Smirnov, *Critical percolation in the plane*, C. R. Acad. Sci. Paris Sér. I Math. **333** no. 3, 239–244 (2001).
- [23] J. J. H. Simmons, P. Kleban and R. M. Ziff, *Percolation crossing formulas and conformal field theory*, J. Phys. A: Math. Theoret. 40, F771–F784 (2007).
- [24] G. Watts, *A crossing probability for critical percolation in two dimensions*, J. Phys. A: Math. Gen. **29**, L363–L368 (1996).
- [25] D. Zagier, *Traces of singular moduli* Motives, polylogarithms and Hodge theory, Part I (Irvine, CA, 1998), 211–244, Int. Press Lect. Ser., 3, I, Int. Press, Somerville, MA, 2002.
- [26] R. M. Ziff, *On Cardy’s formula for the critical crossing probability in 2D percolation* J. Phys. A 28, 1249–1255 (1995); *Proof of crossing formula for 2D percolation* 6479–6480 (1995).